

Properties of Quasar-Galaxy Associations and Gravitational Mesolensing by Halo Objects

Yu.L.Bukhmastova

St.Petersburg State University, St.Petersburg, Russia
bukh_julia@mail.ru, bukh@astro.spbu.ru

1. INTRODUCTION

The problem of physical associations between distant quasars and nearby galaxies has been discussed in the literature for more than thirty years. Burbidge *et al.* [1] published a catalog of galaxy-quasar associations containing 577 quasars and 500 galaxies. They indicated that quasars show a tendency to be located near the halos of normal galaxies much more often than expected in the case of chance projections, and that this physical relationship requires an explanation. The physical connection between quasars and galaxies was first interpreted as a result of gravitational lensing by halo stars in the galaxies (microlensing) by Canizares [2]. However, in subsequent studies [3-5], it was shown that gravitational microlensing cannot explain the observations, due to the extremely low surface density of weak quasars, which should be amplified by microlensing. In addition, Arp [6] presented several arguments against the gravitational microlensing hypothesis.

(1) Quasars in pairs with nearby galaxies are predominantly separated from the galaxy by several galaxy diameters, so that microlenses should occupy a huge volume of space around the galaxies. The observed number of associations requires anomalously large halo masses.

(2) In groups of galaxies, quasar-galaxy associations are more often encountered for companion galaxies than for dominant galaxies in the group.

(3) Analysis of archival data on the variability of quasars in associations [7] showed a lack of variability on timescales of several dozen years, in contradiction with expectations of the microlensing hypothesis.

As a result, Arp [6] concluded that the redshifts of quasars could have a non-cosmological nature.

Baryshev and Ezova (Bukhmastova) [8] attempted to explain the appearance of quasar-galaxy pairs by theorizing that distant active galactic nuclei experienced mesolensing by globular clusters in the halos of more nearby galaxies. In this case, the quasar corresponds to a distant active galactic nucleus whose brightness is amplified by several magnitudes. Estimates of the expected number of associations based on calculations of the probability that active galactic nuclei will be lensed lead to values of $10^2 - 10^4$, assuming brightness enhancements of $5^m - 8^m$. If the mean brightness enhancement is

lowered 3^m , the number of expected associations grows to 10^5 (based on the number of active galactic nuclei being $\sim 10^4/\text{deg}^2$ for galaxies with apparent magnitudes to 29^m). In light of these hypotheses, the arguments of Arp [6] lose their force.

The current paper is a logical continuation of [8]. We do not yet consider the problem of the luminosity function of quasars [4,5], which will be the focus of the next paper in this series. Here, we compose a new, appreciable expanded, catalog of pairs. The number of pairs selected using the new observational material (8382 pairs) is in agreement with the predictions of theoretical computations. The search for pairs was carried out in such a way that the linear distance between the galaxy and projected quasar does not exceed 150 kpc. We used four criteria for the pair search:

- (1) there are the spatial coordinates α, δ and z of the quasars and galaxies, so that the quasar magnitude m_Q is known;
- (2) the quasar must be located farther than the galaxy (i.e., $z_Q > z_G$);
- (3) the quasar should be projected onto the halo of the galaxy;
- (4) the galaxy should have $z_G > 4 \cdot 10^{-4}$.

The criteria for selecting new pairs were based on the main assumptions of the gravitational lensing hypothesis. In particular, this theory led to criteria (2) and (3). Criterion (4) was derived in [8].

The structure of the paper is as follows. Section 2 describes the algorithm for selecting the pairs. Section 3 is dedicated to a general analysis of the resulting pairs. In section 4, presented histograms reflecting an important property of the associations and attempt to analyze the result obtained. Section 5 is concerned with the Hubble diagram for galaxies and quasars, and Section 6 discusses possible tests for the adopted model. We briefly summarize our main conclusions in Section 7.

2. PAIR SELECTION ALGORITHM

Since the quasars are located at greater distances than the galaxies, we will initially project the quasar onto a celestial sphere with a radius corresponding to the distance to the galaxy. We denote the smallest distance between the quasar and galaxy projections $l = GQ$ (Fig.1). We now solve for the spherical triangle PGQ with sides $PG = 90 - \delta_G$, $PQ = 90 - \delta_Q$, and $GQ = l$. According to the cosine theorem,

$$\cos l = \cos \delta_G \cos \delta_Q \cos(\alpha_Q - \alpha_G) + \sin \delta_G \sin \delta_Q.$$

The linear distance x in kpc is $x = z_G \frac{\sqrt{1 - \cos^2 l}}{\cos l} 5 \cdot 10^6$ for $H = 60 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

Thus, if $x \leq 150\text{kpc}$ and $z_G > 4 \cdot 10^{-4}$, the quasar– galaxy pair is specified.

The input quasar catalog [9] is available at the web address:

<ftp://cdsarc.u-strasbg.fr/cats/VII/207>

The galaxy catalog [10] is available at the address:

<http://www-obs.univ-Lyon1.fr>

Our application of the above simple selection operation resulted in the new catalog of associations, which can be found at the web address:

http://www.astro.spbu.ru/staff/Baryshev/gl_dm.htm

3. MAIN PROPERTIES OF THE CLOSE QUASAR-GALAXY PAIRS

The selection criteria indicated above are satisfied by 77843 galaxies and 11358 quasars. Of these, 1054 galaxies and 3164 quasars were included in pairs, making up a total of 8382 quasar-galaxy pairs. Cases when one quasar is near several galaxies are encountered fairly frequently. Examples of such situations are presented in Table 1.

1	2	3	4	5	6	7	8	9
7627	TEX 1322+479	2.26	20.5	PGC0045939	0.0005	14.2	112.6	0.0002
7628	TEX 1322+479	2.26	20.5	PGC0046039	0.0006	11.7	140.7	0.0003
7629	TEX 1322+479	2.26	20.5	PGC0046127	0.0009	13.5	115.6	0.0004
7630	TEX 1322+479	2.26	20.5	PGC0047270	0.0015	13.9	139.1	0.0007
7631	TEX 1322+479	2.26	20.5	PGC0047404	0.0015	8.3	141.3	0.0007
7632	TEX 1322+479	2.26	20.5	PGC0047413	0.0016	9.9	141.9	0.0007
7880	SBS 1400+541	0.646	17.5	PGC0049448	0.0005	12.4	46.7	0.0007
7881	SBS 1400+541	0.646	17.5	LEDA0140246	0.0006	-	0.7	0.0009
7882	SBS 1400+541	0.646	17.5	PGC0050063	0.0008	7.9	31.1	0.0012
7883	SBS 1400+541	0.646	17.5	PGC0050216	0.0009	10.9	37.5	0.0014
7884	SBS 1400+541	0.646	17.5	PGC0050262	0.0010	13.9	63.2	0.0016
7885	SBS 1400+541	0.646	17.5	LEDA0165626	0.0005	-	15.5	0.0008
7886	SBS 1400+541	0.646	17.5	LEDA0165627	0.0010	-	38.0	0.0015
7887	SBS 1400+541	0.646	17.5	LEDA0165629	0.0010	-	48.7	0.0015
8324	Q 2315-4230	2.83	20.0	LEDA0123614	0.0538	16.1	71.8	0.019
8325	Q 2315-4230	2.83	20.0	PGC0071001	0.0052	10.6	64.9	0.0019
8326	Q 2315-4230	2.83	20.0	PGC0071031	0.0053	11.1	57.2	0.0019
8327	Q 2315-4230	2.83	20.0	PGC0071066	0.0056	10.9	99.8	0.0020

Table 1. Examples from the catalog of quasar-galaxy associations showing several galaxies near a single quasar. The columns contain: (1) pair number in the original catalog, (2) quasar name, (3) quasar redshift, (4) quasar apparent magnitude, (5) galaxy name, (6) galaxy redshift, (7) galaxy apparent magnitude, (8) distance between the galaxy and projection of the quasar onto the plane of the galaxy in kpc, (9) $a = z_G/z_Q$. The table presents data for three quasars and eighteen galaxies with which they are associated.

1	2	3	4	5	6	7	8	9
895	Q 0054+0200	1.872	18.4	PGC0003844	0.0008	9.6	137.6	0.0004
934	Q 0054+0236	1.654	17.6	PGC0003844	0.0008	9.6	139.1	0.0005
948	Q 0055+0141	2.232	18.6	PGC0003844	0.0008	9.6	123.1	0.0003
983	Q 0055+0225	0.373	18.6	PGC0003844	0.0008	9.6	119.6	0.0021
995	UM 294	1.914	17.1	PGC0003844	0.0008	9.6	146.4	0.0004
1059	PC 0056+0125	3.154	18.6	PGC0003844	0.0008	9.6	98.8	0.0002
1060	Q 0056+0009	0.613	18.0	PGC0003844	0.0008	9.6	148.9	0.0013
1066	Q 0056+0118	1.1	18.1	PGC0003844	0.0008	9.6	98.8	0.0007
1095	Q 0057+0000	0.776	17.2	PGC0003844	0.0008	9.6	149.4	0.0010
1106	Q 0057+0230	0.716	18.3	PGC0003844	0.0008	9.6	90.5	0.0011
1147	PHL 938	1.959	17.16	PGC0003844	0.0008	9.6	67.4	0.0004
1149	Q 0058+0215	2.868	18.91	PGC0003844	0.0008	9.6	71.2	0.0003
1171	Q 0058+0218	0.929	17.7	PGC0003844	0.0008	9.6	67.0	0.0008
1182	Q 0058+0121	1.432	17.6	PGC0003844	0.0008	9.6	66.8	0.0005
1193	Q 0058+0205	0.6	18.5	PGC0003844	0.0008	9.6	58.3	0.0013
1195	Q 0059+0147	1.143	18.0	PGC0003844	0.0008	9.6	55.7	0.0007
1255	Q 0059+0035	2.545	18.5	PGC0003844	0.0008	9.6	95.1	0.0003
1311	Q 0100+0228	1.543	18.4	PGC0003844	0.0008	9.6	50.8	0.0005
1314	Q 0100+0146	1.909	18.5	PGC0003844	0.0008	9.6	30.3	0.0004
1315	UM 301	0.393	16.39	PGC0003844	0.0008	9.6	32.0	0.0020
1330	Q 0100+0106	1.405	18.5	PGC0003844	0.0008	9.6	57.3	0.0005
1335	Q 0101+0024	1.436	18.9	PGC0003844	0.0008	9.6	100.6	0.0005
1344	Q 0101+0009	0.394	17.4	PGC0003844	0.0008	9.6	116.6	0.0020
1370	Q 0101+0118	1.133	18.61	PGC0003844	0.0008	9.6	38.3	0.0007
1403	Q 0102+0036	0.649	18.4	PGC0003844	0.0008	9.6	84.3	0.0012
1408	Q 0102+0241	1.509	18.6	PGC0003844	0.0008	9.6	55.3	0.0005
1414	LMA 15	2.7	0.0	PGC0003844	0.0008	9.6	6.7	0.0003
1444	Q 0103+0234	1.7	18.0	PGC0003844	0.0008	9.6	49.8	0.0005
1448	Q 0103+0024	1.075	17.4	PGC0003844	0.0008	9.6	99.7	0.0007
1454	PC 0103+0123	3.066	19.7	PGC0003844	0.0008	9.6	36.4	0.0003
1467	Q 0103+0123	0.782	18.8	PGC0003844	0.0008	9.6	37.9	0.0010
1475	BRI 0103+0032	4.437	18.6	PGC0003844	0.0008	9.6	92.5	0.0002
1486	Q 0103-0014	1.629	18.5	PGC0003844	0.0008	9.6	144.7	0.0005
1487	Q 0104+0001	0.91	18.3	PGC0003844	0.0008	9.6	127.6	0.0008
1493	Q 0104+0030	1.874	18.5	PGC0003844	0.0008	9.6	96.9	0.0004
1499	PC 0104+0215	4.171	19.73	PGC0003844	0.0008	9.6	41.4	0.0002
1578	PKS 0106+01	2.107	18.39	PGC0003844	0.0008	9.6	73.0	0.0004
1606	Q 0107+0051	0.966	19.0	PGC0003844	0.0008	9.6	106.4	0.0008
1613	Q 0107+0022	1.968	18.3	PGC0003844	0.0008	9.6	131.3	0.0004
1640	MS 01080+0139	0.713	17.49	PGC0003844	0.0008	9.6	96.3	0.0011
1643	Q 0108+0028	2.005	18.25	PGC0003844	0.0008	9.6	134.2	0.0004
1651	Q 0108+0030	0.428	19.0	PGC0003844	0.0008	9.6	135.8	0.0018
1678	MS 01094+0242	0.262	18.1	PGC0003844	0.0008	9.6	132.0	0.0029
1682	UM 87	2.343	17.3	PGC0003844	0.0008	9.6	126.5	0.0003
1705	PB 6325	0.774	17.8	PGC0003844	0.0008	9.6	146.0	0.0010
1708	PB 6327	1.509	18.1	PGC0003844	0.0008	9.6	149.5	0.0005

Table 2. Examples from the catalog of quasar-galaxy associations showing a number of quasars near a single galaxy. The columns contain: (1) pair number in the original catalog, (2) quasar name, (3) quasar redshift, (4) quasar apparent magnitude, (5) galaxy name, (6) galaxy redshift, (7) galaxy apparent magnitude, (8) distance between the galaxy and projection of the quasar onto the plane of the galaxy in kpc, (9) $a = z_G/z_Q$. The table presents data for the galaxy PGC 0003844 and 47 quasars with which it is associated.

One simple explanation for the data can be obtained if the distribution of galaxies along the line of sight has a fractal nature; i.e., if, in accordance with [11], we adopt the following parametric representation for the concentration of objects along the line of sight between the source and observer:

$$n_l(R) = 0.5n_{ol}\left[\left(\frac{R}{R_0}\right)^{D_F-3} + \left(\frac{R_s - R}{R_0}\right)^{D_F-3}\right],$$

where $n_{ol}(R_0)$ is the concentration of objects at the distance R_0 , R is the distance to the lens, R_s is the distance to the source, and D_F is the fractal dimension of the galaxy distribution. Usually, a uniform distribution of lenses is obtained if $D_F = 3$, in which case $n_l(R) = \text{const.}$ The observed value for the fractal dimension is close to $D_F \sim 2$ [12,13]. The character of this relation is such that, if some quasar (according to the Barnotti–Tyson hypothesis, the nucleus of an active galaxy) is projected onto the halo of a nearby galaxy, the probability that the quasar is also projected onto other galaxies is enhanced, since the galaxy is part of a fractal structure. The inverse is also true, since two objects of a single class along the line of sight in a fractal structure have equal validity relative to the path of a light ray. This means that cases when there are several quasars near a single galaxy should be rare. Examples of such cases are presented in Table 2. The galaxies PGC 0003290, PGC 0003589, PGC 0001014, PGC 0003238, PGC 0003721, PGC 0002789, and a number of others are surrounded by tens or hundreds of quasars.

Why should a distant quasar be projected onto the halo of a more nearby galaxy? There exist at least three possible explanations.

(1) If the galaxy is very close to the observer, it covers a relatively large fraction of sky. There is some probability for the chance projection of quasar onto the halo of the nearby galaxy.

(2) If the spatial distribution of galaxies is fractal and at least some fraction of quasars are the nuclei of active galaxies (i.e., they belong to the same general class of objects—galaxies), there should be a significant number of pairs due to the properties of fractal distribution, as noted above.

(3) Due to gravitational mesolensing by halo objects with intermediate mass, such as globular clusters, whose number in galactic halos can reach several thousand [4], some fraction of objects viewed through these lenses will appear to have enhanced luminosities and will be interpreted as being quasars [3]. As a result, the clustering of the mass in the halos of nearby galaxies “magnifies” the light from a distant galaxy that would otherwise simply not be visible. The result is the detection of a close quasar–galaxy pair. The result is the detection of a close quasar–galaxy pair. The appearance of numerous quasars around a single galaxy can be explained by the large number of globular clusters acting as lenses in the halo.

We will analyze the distribution of the galaxies along the line of sight from the observer to the quasar in order to elucidate the origin of the observed associations.

4. MUTUAL LOCATIONS OF THE GALAXIES AND QUASARS IN ASSOCIATIONS

The distance to the quasar (which we take to be indicated by its redshift) is denoted z_Q . The analogous distance to the galaxy is z_G . We introduce the quantity $a = z_G/z_Q$, which is the normalized distance from the observer to the galaxy in a system in which the distance to the quasar is equal to unity. We determined a for all the quasar-galaxy pairs. The result is shown as a histogram of x as a function of number of pairs with that a value (not to scale). We obtain four relations for various halo sizes (50 and 150 kpc) and quasar redshifts (Fig.2).

Even without a more careful analysis, we can see that galaxies in associations are preferably located either near the observer, $a < 0.1$, or near the quasar, $a > 0.9$, avoiding intermediate distances between the observer and quasar. The appearance of the first "tail" at $a < 0.1$ is not surprising. Most galaxies are located nearby, so that the substantial contribution to the number of pairs from such galaxies could be an observational selection effect. What about the second "tail" at $a > 0.9$, which is stable to variation of the halo size and quasar redshift? Similar dependences were obtained for the catalog [1] in [8], based on 241 pairs. The data have now been expanded appreciably to 8382 pairs, but the dependences have remained essentially unchanged. Let us try to elucidate their intrinsic properties. We assume that the following effects are responsible for these dependences to some degree:

- (1) the fractal distribution of galaxies along the line of sight, with a fractal dimension close to 2;
- (2) gravitational lensing by globular clusters in the galactic halos (or by other halo objects displaying clustering with a King mass distribution);
- (3) chance projected pairs.

We studied this question using computer simulations. We fixed the galaxy positions and scattered the quasars randomly, with z_Q from 0.1 to 3. In this way, we attempted to remove the effect of possible lensing. There remain to effects that could lead to the formation of pairs: random projections and nature of the distribution of the galaxies along the line of sight. Figure 3 shows the result of identifying pairs for the case of a random distribution of background quasars. The results have changed both qualitatively and quantitatively. The number of pairs has dropped dramatically, to 2000

(compared to 8382), and the second tail in the distributions has disappeared; i.e., its presence in the observational data is not due to random effects.

As a second step in our computer simulations, we randomly specified the positions of 77483 galaxies, with z_G from 0 to 0.25, and of 11358 quasars, with z_Q from 0.1 to 3. Figure 4 shows the result of identifying pairs for this case of randomly distributed galaxies and quasars. An analysis of this histogram indicates that about 1200 pairs could be obtained due to random positional coincidences. However, another qualitative result is more reliable and interesting: in the case of chance coincidences, the quasars in pairs are not projected onto nearby galaxies whose distances are close to those of the quasars; i.e., again, the tail at $a > 0.9$ disappears.

Thus, crude estimates of the model relations suggest that about 15% of the pairs formed due to random projected positional coincidences. In the framework of our adopted assumptions, the remaining pairs form as a result of the fractal distribution of galaxies along the line of sight and due to lensing by halo objects with a King mass distribution, with the effects of lensing and of the fractal galaxy distribution complementing each other [8]. Let us now consider the following selection of 8382 quasar-galaxy pairs. If some quasar is projected onto the halos of several galaxies, we select the pair with the minimum distance between the quasar and galaxy projected onto a single galaxy, we consider all these pairs, since several hundred, or even several thousand, globular clusters can be located in the galaxy's halo, each of which can act as a lens, resulting in several amplified active nuclei of distant galaxies, interpreted as quasars. We now construct the same histogram for such a sample of 3164 quasars and 1054 galaxies, shown in Fig.5. Qualitatively, the results are as before: there are excesses of pairs with $a < 0.1$ and $a > 0.9$. Note the numerical values for $a > 0.9$ ("tail" with $a < 0.1$ has a high fraction of chance pairs, and is not informative in a numerical sense). It turns out that the postulated lenses are located primarily in halos less than 50 kpc in size. The histogram shows that about 83% of the lenses are located in such halos, whose size does not exceed 50 kpc while the remaining 17% of the lenses are located at distances of 50 to 150 kpc. If these lenses are globular clusters, they should be located at precisely such distances from the centers of their galaxies, in accordance with the data of [14].

As an example, we present data for 146 globular clusters in the Milky Way (Fig.6). We can see that 140 of the 146 globular clusters are located in the halo out to distances of 50 kpc, in agreement with the histograms in Figs.2 and 5. The same dependences of the number of globular clusters with distance from the galactic center are characteristic, for example, A754, A1644, A2124, A2147, A2151, A2152 [15], as well as for NGC4874, NGC4889, NGC4472, NGC4486 [16].

5. THE HUBBLE DIAGRAM $z = f(m)$ FOR GALAXIES AND QUASARS

Figure 7 presents a Hubble diagram $z = f(m)$, for the 77483 LEDA galaxies and 11358 quasars. All the quasars are shifted to the left from the straight line corresponding to the "0.2m" law.

In the context of the gravitational lensing hypothesis, this shift for the quasars can be explained as the effect of brightness amplification during lensing. If the quasars are the amplified nuclei of active galaxies, they can be shifted to the strip corresponding to the brightest galaxies by "removing" several magnitudes from their brightnesses. The presence of a distinct lower boundary to the region occupied by the quasars can be explained by the fact that there are objects brighter than -23^m in the initial catalog of quasars.

6. OBSERVATIONAL TESTS

Since the observational data for the quasar-galaxy associations are consistent with the gravitational lensing model, this raises the question of further observational tests of the model. We propose the following observational tests of the properties of the quasars that are members of associations.

(1) The expected angular separation of multiple images of quasars due to lensing by objects such as globular clusters is several milliarcseconds. Therefore, it is important to study the structure of the compact radio components of quasars in associations that have sufficient radio fluxes for VLBI observations. Of course, the theory does not exclude the possibility of a single image being formed. In addition, the image splitting could be unobserved due to insufficient dynamical range for the observations.

(2) The expected variability of the quasar's brightness due to the motion of the globular clusters takes place on time scales of more than a thousand years. Variability on timescales of less than a year is also possible, as a consequence of microlensing by individual globular cluster stars.

(3) Since globular clusters are located in the halos of galaxies, absorption lines with z_{abs} corresponding to the redshift of the galaxy may arise in the quasar spectra. It would be interesting to continue the work begun in [17], comparing the properties of quasars with absorption lines and quasars in associations. In particular, the well known increase in the number of absorption lines with approach toward the quasar could be associated with the fractal nature of the large-scale distribution of galaxies along the line of sight. In this case, we should also expect an increase in the number of absorption lines with approach toward the observer.

(4) Comparisons of the spectral properties of quasars in associations with those of various types of active galactic nuclei could serve as a probe of

the structure of the emission-line formation region, which could be affected differently during lensing by King objects.

(5) Analysing the properties of quasar-galaxy associations, we can estimate the fraction of galaxies of various morphological types in the total number of associations and the distribution of globular clusters in each type of galaxy.

(6) In the case of quasar with jets, the curvature in the jet trajectories that is not associated with real curved motion of the material ejected from the core, but is related to the refraction of rays in the presumed lens is possible.

(7) In the hypothesis of gravitational microlensing of quasars, the Hubble diagram presented above can be used to estimate the amplification coefficient for quasars, by moving each quasar to the strip of brightest galaxies. Knowing the dependence of the amplification coefficient on the position of the lens between the observer and source, it is possible to estimate the distance to the postulated lens.

(8) It is possible to devise a computer simulations of the large-scale fractal distribution of galaxies and quasars, in order to organize searches for quasar-galaxy pairs and compare the resulting data with the catalog of observed associations.

7. CONCLUSION

We have derived a new catalog of close quasar– galaxy pairs including 1054 galaxies and 3164 quasars comprising 8382 pairs. All the galaxies in the associations show a strong tendency to be located close to either the observer or the quasar, avoiding intermediate positions along the line of sight to the quasar. This property of the pairs and their considerable number can easily be explained if we adopt the following assumptions.

1) The quasars in associations could be gravitationally amplified active nuclei of distant galaxies. There are sufficient numbers of active galactic nuclei to provide the observed number of quasars if there is a probability of 10^{-4} for a brightness amplification by 3^m .

2) Galaxies on scales ~ 150 Mpc have a fractal distribution with a fractal dimension close to two. A uniform galaxy distribution makes it difficult to explain the enhanced number of pairs with $a > 0.9$.

3) The role of the gravitational lenses can be played by objects such as globular clusters or clusters of dark matter characterized by a King density distribution, located primarily in galactic halos at distances of up to 50 kpc from the galactic center. The importance of the King distribution is that it has a conical caustic that can explain the enhanced probability for the galaxies harboring the lenses to be near either the observer or the quasar.

The point and isothermal-sphere lens models that are often considered in the literature give an enhanced probability for the galaxy to be located in a central position between the observer and the source, in contradiction with the observed properties of the association. The derived histograms indicate that quasars in associations are most often projected onto the galactic halo at distances out to 50 kpc, in spite of the fact that the pair selection criteria allowed projections out to 150 kpc. This provides an additional argument that the relationship between the distant quasars and nearby galaxies is associated with globular clusters in the galaxy halos.

4) Quasars that were not included in the catalog of associations can also be gravitationally amplified galactic nuclei, since it is probable that a galaxy surrounded by lensing objects is present near the projection of the quasar and is simply not detected. This possibility follows from the above histograms.

The hypothesis that we are dealing with gravitational lensing of the distant nuclei of galaxies can provide a simple physical interpretation of the Arp effect; i.e., the observed frequency with which quasar-galaxy associations are encountered. The quasar redshifts are cosmological in nature, and do not require any new physics.

ACKNOWLEDGMENTS

The author is grateful to Yu.V.Baryshev for constructive discussions of the problem, and also to D.S. Bukhmastov for appreciable technical help in preparing the manuscript for publication.

REFERENCES

1. G.Burbidge, A.Hewitt, J.V.Narlikar, and P.Das Gupta, *Astrophys. J., Suppl. Ser.* **74**, 675 (1990).
2. C.R.Canizares, *Nature* **291**, 620 (1981).
3. M.Vietri and J.P.Ostriker, *Astrophys. J.* **267**, 488 (1983).
4. E.V.Linder and P.Schneider, *Astron. Astrophys.* **204**, L8 (1988).
5. P.Schneider, J. Ehlers, and E.E.Falko, *Gravitational Lenses* (Springer-Verlag, New York, 1992).
6. H.Arp, *Quasar, Redshifts and Controversies* (Cambridge Univ.Press,Cambridge, 1988).
7. W.C.Keel, *Astrophys.J.* **259**, L1 (1982).
8. Yu.V.Baryshev and Yu.L.Ezova, *Astron.Zh.* **74**, 497 (1997) [*Astron.Rep.* **41**,436(1997)].
9. M.P. Veron-Cetty and P.Veron, *A Catalogue of Quasars and Active Galactic Nuclei* (European Southern Observatory, Garching,1998),

- ESO Scientific Report Series, Vol. 18.
10. G.Paturel, H.Andermach, L.Bottinelli, et al.,
*Astron.Astrophys.,Suppl.Ser.***124**, 109 (1997).
 11. Yu.V.Baryshev, A.A.Raikov, and A.A.Tron, in *Gravitational Lenses in the Universe*, 31st Liege International Astrophysics Colloquium, 1993, p.365.
 12. Yu.V.Baryshev, F.Sylos-Labini, M.Montuori and L.Pietronero,
*Vistas Astron***38**, 419 (1994).
 13. F.Sylos-Labini, M.Montuori and L.Pietronero, *Phys. Rep.*
293, 61 (1998).
 14. W.E.Harris, *Astron. J.* **112**, 1487 (1996);
<http://www.physics.mcmaster.ca/Globular.html>.
 15. J.P.Blakeslee, <http://babbage.sissa.it/astro-ph/9906356>.
 16. J.P.Blakeslee and J.L.Tonry, *Astrophys.J.* **442**, 579 (1995).
 17. A.F.Dravskikh, *Astron.Zh.* bf 73, 19 (1996) [*Astron.Rep.* **40**, 13 (1996)].

This figure "fig1.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>

This figure "fig2.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>

This figure "fig3.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>

This figure "Fig4.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>

This figure "fig5.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>

This figure "fig6.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>

This figure "fig7.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/0206456v1>